

UMN-TH-1331/95  
hep-th/9505113  
June 1, 1995 (corrected)  
To appear in Phys. Lett. B

# Spontaneous Magnetization in Lorentz Invariant Theories

Denne Wesolowski and Yutaka Hosotani

*School of Physics and Astronomy, University of Minnesota  
Minneapolis, Minnesota 55455, U.S.A.*

wesolowski@mnhep.hep.umn.edu

yutaka@mnhep.hep.umn.edu

## Abstract

In a class of three-dimensional Abelian gauge theories with both light and heavy fermions, heavy chiral fermions can trigger dynamical generation of a magnetic field, leading to the spontaneous breaking of the Lorentz invariance. Finite masses of light fermions tend to restore the Lorentz invariance.

It has been shown in refs. [1, 2] that in a variety of classes of three-dimensional gauge theories described by

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{\kappa_0}{2} \varepsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho \\ & + \sum_a \frac{1}{2} [\bar{\psi}_a, (\gamma_a^\mu (i\partial_\mu + q_a A_\mu) - m_a) \psi_a] , \end{aligned} \quad (1)$$

the Lorentz invariance of the vacuum is spontaneously broken via dynamical generation of a magnetic field. In other words, it has been established that for these models there exists a variational ground state, in which  $\langle F_{12}(x) \rangle = -B \neq 0$ , that has a lower energy density than the perturbative ground state. Quantum fluctuations play a crucial role in lowering the energy density for these states.<sup>1</sup> In the previous papers models with  $\kappa_0 \neq 0$  and  $m_a \ll q_a^2$  were considered.

There are two crucial ingredients. First, once a magnetic field is dynamically generated, the fermion energy spectrum is characterized by that of Landau levels. The lowest level has an energy  $\epsilon_{a0} = m_a$ . In the massless fermion limit  $\epsilon_{a0} = 0$  so that a perturbative ground state is infinitely degenerate. This degeneracy is lifted by interactions. With certain conditions satisfied, a variational state, in which the lowest Landau levels of chirality + fermions are completely occupied, whereas those of chirality - fermions are empty, has the lowest energy.[1] Secondly it is recognized that the lowering of the energy density by a non-vanishing magnetic field is mainly due to the shift in zero-point energies of photons. If  $\kappa_0 \neq 0$ , a photon is topologically massive in perturbation theory. Once  $B \neq 0$  is generated, a photon becomes a gapless mode, serving as the Nambu-Goldstone boson associated with the spontaneous breaking of the Lorentz invariance. Hence zero-point energies are shifted, resulting in lowering the energy density of the ground state.[2, 4]

Recently Gusynin, Miransky, and Shovkovy have shown that in a model with an external magnetic field and an additional four-fermi interaction, flavor-nonsinglet condensate  $\langle \bar{\psi}_a \psi_a \rangle$  can be spontaneously generated.[5, 6] Again the existence of infinitely degenerate perturbative ground states in the massless fermion limit plays a crucial role.

We here stay within renormalizable theories described by (1) with general fermion

---

<sup>1</sup>Cea has considered a variational ground state with  $B \neq 0$ , which, however, differs from ours.[3]

content to obtain a criterion for the spontaneous magnetization ( $B \neq 0$ ) to occur. We shall also evaluate contributions to the energy density due to finite fermion masses.

A brief review is in order. In (1)  $\psi_a$  is a two-component Dirac spinor. These spinors, in  $2 + 1$  dimensions, are characterized by the signature of the two-dimensional Dirac matrices via the chirality  $\eta_a = \frac{i}{2} \text{Tr} \gamma_a^0 \gamma_a^1 \gamma_a^2 = \pm 1$ . We utilize the representation  $\gamma_a^\mu = (\eta_a \sigma_3, i\sigma_1, i\sigma_2)$ . Since the model is invariant under charge conjugation and the transformation  $m_a \rightarrow -m_a$  is equivalent to the transformation  $\gamma_a^\mu \rightarrow -\gamma_a^\mu$  or  $\eta_a \rightarrow -\eta_a$ , one can take  $q_a > 0$  and  $m_a \geq 0$  without loss of generality.

As in [1] we have the expression for the energy density

$$\begin{aligned}
\Delta \mathcal{E} &= \mathcal{E}(B \neq 0) - \mathcal{E}(B = 0) \\
&= \frac{1}{2} B^2 + \Delta \mathcal{E}_{\text{f.z.e.}} + \Delta \mathcal{E}_{\text{fluct}} \\
\Delta \mathcal{E}_{\text{fluct}} &\sim \Delta \mathcal{E}_{\text{RPA}} \\
&= -\frac{i}{2} \int \frac{d^3 p}{(2\pi)^3} \left\{ \text{tr} \ln (1 - \Gamma^{(2)} D_0) \Big|_{qB, \nu} - (B \rightarrow 0) \right\} \\
&= -\frac{i}{2} \int \frac{d^3 p}{(2\pi)^3} \ln \frac{(1 + \Pi_0) \left\{ 1 + \frac{1}{p^2} (p_0^2 \Pi_0 - \vec{p}^2 \Pi_2) \right\} - \frac{1}{p^2} (\kappa_0 - \Pi_1)^2}{(B \rightarrow 0)} . \quad (2)
\end{aligned}$$

$\Delta \mathcal{E}_{\text{f.z.e.}}$  is the shift in fermion zero-point energies in a given uniform magnetic field.  $\Delta \mathcal{E}_{\text{fluct}}$  represents the shift in the energy density due to quantum fluctuations of fields, and is approximated by the RPA correction  $\Delta \mathcal{E}_{\text{RPA}}$ . The vertex functions  $\Gamma$ , the bare and full photon propagators  $D_0$  and  $D$ , and the invariant functions  $\Pi$ 's are related by

$$\begin{aligned}
\Gamma^{\mu\nu}(p) &= (D_0^{-1} - D^{-1})^{\mu\nu} \\
&= (p^\mu p^\nu - p^2 g^{\mu\nu}) \Pi_0 + i \epsilon^{\mu\nu\rho} p_\rho \Pi_1 \\
&\quad + (1 - \delta^{\mu 0})(1 - \delta^{\nu 0})(p^\mu p^\nu - \vec{p}^2 \delta^{\mu\nu})(\Pi_2 - \Pi_0). \quad (3)
\end{aligned}$$

The superscript for  $\Gamma$  in (2) indicates that  $\Gamma(p)$  is approximated by  $\mathcal{O}(e^2)$  diagrams.

Much as in [1, 2] we obtain a consistency condition which the fermion content must satisfy in order that there be symmetry breaking. Equations of motion imply, in the true vacuum, that  $\kappa_0 \langle F_{12} \rangle = \langle J^0 \rangle$ . Since  $\langle J^0 \rangle = \Pi_1(0) \langle F_{12} \rangle$ , the relation  $\kappa_0 = \Pi_1(0)$  must be satisfied in order for  $B = -\langle F_{12} \rangle$  to be nonvanishing. This consistency condition

was shown to relate to the Nambu-Goldstone theorem associated with the spontaneous breaking of the Lorentz invariance. The dynamically generated nonzero magnetic field must accompany a gapless excitation, which is the photon of the theory.<sup>2</sup> Since the induced Chern-Simons term has a coefficient  $-\Pi_1(0)$ , one can phrase that the bare Chern-Simons term must be exactly cancelled by the induced Chern-Simons term in order to have  $B \neq 0$ .

We now consider some definite cases of fermion content. Suppose that there are light fermions ( $m_{\text{light}} \ll q^2$ ) and heavy fermions ( $m_{\text{heavy}} \gg q^2$ ). In perturbation theory

$$\begin{aligned} \Pi_0|_{B=0} &= \Pi_2|_{B=0} = \sum \frac{q_a^2}{8\pi} \frac{1}{(-p^2)^{1/2}} \left( \sqrt{z_a} + (1 - z_a) \sin^{-1} \frac{1}{\sqrt{1 + z_a}} \right) \\ \Pi_1(p)|_{B=0} &= - \sum \eta_a \frac{q_a^2}{4\pi} \sqrt{z_a} \sin^{-1} \frac{1}{\sqrt{1 + z_a}}, \quad z_a = \frac{4m_a^2}{-p^2}. \end{aligned} \quad (4)$$

This gives, for the part of heavy fermion contributions,[11, 12, 13]

$$\Pi_1^{\text{heavy}}(0)|_{B=0} = - \sum_{\text{heavy}} \eta_a \frac{q_a^2}{4\pi} = \kappa_{\text{heavy}}. \quad (5)$$

In the presence of  $B \neq 0$  the energy spectrum of fermions has the structure of Landau levels  $\epsilon_{an} = \sqrt{m_a^2 + 2nq_a|B|}$  ( $n = 0, 1, 2, \dots$ ). There is asymmetry in the  $n = 0$  modes.  $\Pi_1(0)$  is found to be

$$\Pi_1(0)|_{B \neq 0} = \frac{1}{2\pi} \sum \eta_a q_a^2 \left( \nu_a - \frac{1}{2} \right). \quad (6)$$

Here  $\nu_a$  is the filling fraction of the Lowest Landau level. We suppose that either  $\nu_a = 0$  (empty) or  $\nu_a = 1$  (completely filled).

The lowest Landau level has an energy  $m_a$ . Hence for heavy fermions we may assume that the lowest Landau levels are completely empty, and assign a filling factor  $\nu_a = 0$ . The approximation is valid for  $|B| \ll m_{\text{heavy}}^{3/2}$ . Comparing (6) with (5), one finds

$$\Pi_1^{\text{heavy}}(0)|_{B \neq 0} = \Pi_1^{\text{heavy}}(0)|_{B=0} \equiv -\kappa_{\text{heavy}} \quad (7)$$

Further heavy fermion contributions to  $\Pi_0$  and  $\Pi_2$  at low energies ( $|p^2| \ll m_{\text{heavy}}^2$ ) are  $O(q^2/m_{\text{heavy}})$ . In other words, the only effect of heavy fermions at low energies is to shift the bare Chern-Simons coefficient  $\kappa_0$  to  $\kappa_0 + \kappa_{\text{heavy}}$ .

---

<sup>2</sup>The possibility of regarding a photon as a Nambu-Goldstone boson has been discussed by Bjorken[8], by Nambu[9], and by Kovner and Rosenstein[10]. Our picture is different from theirs. In our case the Lorentz invariance is spontaneously broken in the physical content.

The variational ground state (true vacuum) is specified with  $\{\nu_a ; a \in \text{light fermions}\}$ . The consistency condition to be satisfied is thus

$$\kappa_0 + \kappa_{\text{heavy}} = \frac{1}{2\pi} \sum_{\text{light}} \eta_a q_a^2 (\nu_a - \frac{1}{2}) . \quad (8)$$

Even if the bare Chern-Simons coefficient  $\kappa_0 = 0$ , the requisite consistency condition can be satisfied.

To be more definite, we consider the following models:

[Model A] The light fermion content is chirally symmetric. In other words, light fermions come in a pair of  $(m_a, q_a, \eta_a = \pm)$ .  $\{\nu_a\}$  must satisfy (8).

[Model B] This is a special case of model A, in which  $\nu_a = 1$  or  $0$  for  $\eta_a = +$  or  $-$ , respectively. In this case the consistency condition is reduced to

$$\kappa_0 + \kappa_{\text{heavy}} = \frac{1}{4\pi} \sum_{\text{light}} q_a^2 . \quad (9)$$

[Model C] In model B we suppose that  $q_a = q$  and  $m_a = m$  ( $a = 1 \sim N_f$ ). The consistency condition is

$$\kappa_0 + \kappa_{\text{heavy}} = \frac{N_f q^2}{4\pi} . \quad (10)$$

The shift in zero-point energies of photons between the perturbative vacuum and the variational ground state is given by [2]

$$\begin{aligned} \Delta\mathcal{E} &= \int \frac{d\vec{p}}{(2\pi)^2} \frac{1}{2} \{ \omega(\vec{p})_{B \neq 0} - \omega(\vec{p})_{B=0} \} \\ &\sim \int \frac{d\vec{p}}{(2\pi)^2} \frac{1}{2} \left\{ \sqrt{\vec{p}^2 + \kappa_{\text{tot}}^2(p)_{B \neq 0}} - \sqrt{\vec{p}^2 + \kappa_{\text{tot}}^2(p)_{B=0}} \right\} , \end{aligned} \quad (11)$$

where  $\kappa_{\text{tot}}(p) = \kappa_0 - \Pi_1(p)$ . In model A  $\Pi_1(p)^{\text{light}} = 0$  identically to  $\mathcal{O}(e^2)$ . Hence, for  $p^2 \ll m_{\text{heavy}}^2$ ,  $\kappa_{\text{tot}}(p)_{B=0} \sim \kappa_0 + \kappa_{\text{heavy}}$ . On the other hand the consistency condition implies that  $\kappa_{\text{tot}}(0)_{B \neq 0} = 0$ . For large  $|\vec{p}|$ ,  $\kappa_{\text{tot}}(p)_{B \neq 0} \sim \kappa_{\text{tot}}(p)_{B=0}$ . The deviation develops below  $l_{\text{ave}}^{-1}$  where  $l_{\text{ave}}$  is a characteristic magnetic length. For  $m_{\text{light}} = 0$  [2]

$$\begin{aligned} \frac{1}{l_{\text{ave}}^2} &= \frac{|\sum_{\text{light}} \eta_a \nu_a q_a^3|}{|\sum_{\text{light}} \eta_a \nu_a q_a^2|} \times |B| && (\text{model A}) \\ &= \frac{\sum_{\text{light}} q_a^3}{\sum_{\text{light}} q_a^2} \times |B| && (\text{model B}) . \end{aligned} \quad (12)$$

To get an estimate of (11), we approximate  $\kappa_{\text{tot}}(p)_{B \neq 0} = 0$  and  $\kappa_{\text{tot}}(p)_{B=0} = \kappa_0 + \kappa_{\text{heavy}}$  for  $|\vec{p}| \leq l_{\text{ave}}^{-1}$ , whereas  $\kappa_{\text{tot}}(p)_{B \neq 0} = \kappa_{\text{tot}}(p)_{B=0}$  for  $|\vec{p}| \geq l_{\text{ave}}^{-1}$ . Then

$$\begin{aligned} \Delta \mathcal{E} &\sim -\frac{1}{8\pi} \frac{|\kappa_0 + \kappa_{\text{heavy}}|}{l_{\text{ave}}^2} + O(l_{\text{ave}}^{-3}) \\ &= -\frac{1}{32\pi^2} \sum_{\text{light}} q_a^3 \cdot |B| \quad (\text{model B}) \end{aligned} \quad (13)$$

for  $|\kappa_0 + \kappa_{\text{heavy}}| l_{\text{ave}} \gg 1$ . As (11) and (13) are negative definite, the energy density  $\mathcal{E}$  is indeed lowered for the vacuum state with  $B \neq 0$  provided that  $\kappa_0 + \kappa_{\text{heavy}}$ ,  $\sum_{\text{light}} \eta_a \nu_a q_a^2$ , and  $\sum_{\text{light}} \eta_a \nu_a q_a^3$  are all non-vanishing and that the consistency condition (8) is satisfied. In particular, even if the bare Chern-Simons term is absent ( $\kappa_0 = 0$ ), heavy fermions can induce a dynamical magnetic field  $B \neq 0$ .

The same conclusion is obtained more convincingly by evaluating (2). We recall [1]

$$\begin{aligned} \Pi_0(p)|_{B \neq 0} - \Pi_0(p)|_{B=0} &= \sum_{\text{light}} \nu_a \frac{q_a^2}{2\pi} \frac{1}{p_0} \left\{ \frac{1}{2 - p^2 l_a^2 - 2m_a p_0 l_a^2} - \frac{1}{2 - p^2 l_a^2 + 2m_a p_0 l_a^2} \right\} + O(B^2) \\ \Pi_1(p)|_{B \neq 0} - \Pi_1(p)|_{B=0} &= \sum_{\text{light}} \eta_a \nu_a \frac{q_a^2}{2\pi} \left\{ \frac{1}{2 - p^2 l_a^2 - 2m_a p_0 l_a^2} + \frac{1}{2 - p^2 l_a^2 + 2m_a p_0 l_a^2} \right\} + O(B^2) \\ \Pi_2(p)|_{B \neq 0} - \Pi_2(p)|_{B=0} &= O(B^2) \end{aligned} \quad (14)$$

where  $l_a^{-2} = |q_a B|$ . When  $m_a = 0$ ,  $\Pi_0(p)_{B=0} = \Pi_2(p)_{B=0} = (\sum_{\text{light}} q_a^2/16)(-p^2)^{-1/2}$  and  $\Pi_1(p)_{B=0} = -\kappa_{\text{heavy}}$  for  $|-p^2| \ll m_{\text{heavy}}^2$ . Inserting these with (14) into (11) gives the result [2]

$$\begin{aligned} \Delta \mathcal{E}_{\text{RPA}} &= -\frac{\sum_{\text{light}} \eta_a \nu_a q_a^3}{2\pi^3} \cdot \tan^{-1} \frac{8 \sum_{\text{light}} \eta_a \nu_a q_a^2}{\pi \sum_{\text{light}} q_a^2} \cdot |B| + O(|B|^{3/2}) \\ &= -\frac{1}{4\pi^3} \tan^{-1} \frac{4}{\pi} \sum_{\text{light}} q_a^3 \cdot |B| + O(|B|^{3/2}) \quad (\text{model B}). \end{aligned} \quad (15)$$

Comparing this with (13), one finds that the shift in zero-point energies accounts for half of the RPA effect.

Recently Cangemi, D'Hoker and Dunne have evaluated the effective potential in a magnetic field, supposing that all Landau levels are empty in the vacuum.[7] They found

that a uniform constant magnetic field configuration shows instability against forming inhomogeneity. Cangemi et al.'s variational vacuum state corresponds to  $\{\nu_a = 0\}$  whereas in our variational vacuum  $\nu_a = 1$  for  $\eta_a = +$ . It is desirable to evaluate the effective potential for a general vacuum state specified with  $\{\nu_a\}$ .

In passing, if  $\nu_a = 0$  in model A, the consistency condition (8) becomes  $\kappa_0 + \kappa_{\text{heavy}} = -(4\pi)^{-1} \sum \eta_a q_a^2 = 0$ . Further, from the symmetry  $\Pi_1(p)_{\text{light}} = 0$ . Hence both  $\kappa_{\text{tot}}(p)_{B=0}$  and  $\kappa_{\text{tot}}(p)_{B \neq 0}$  vanish for  $p^2 \ll m_{\text{heavy}}^2$ . In other words the state with  $(B \neq 0, \nu_a = 0)$  has a higher energy density than the state with  $B = 0$ .

It is straightforward to evaluate  $O(m_a)$  corrections to  $\Delta\mathcal{E}$ . We evaluate them in model C, supposing  $q_a = q, m_a = m \neq 0$ .  $\Delta\mathcal{E}/q^6$  is a function of two dimensionless parameters  $m/q^2$  and  $|B|/q^3$ . We shall find an expression for  $\Delta\mathcal{E}$  valid for  $m^2/q^4 \ll |B|/q^3 \ll 1$ . (Note  $(ml)^2 = m^2/(q|B|) \ll 1$ .)

Corrections to  $\Delta\mathcal{E}_{\text{RPA}}$  are found from (2) and (14).  $\Pi_0$  and  $\Pi_1$  give corrections of  $O(m)$  and  $O(m^2)$ , respectively. Hence a dominant correction comes from  $\Pi_0$  and is found to be

$$\begin{aligned} \Delta\mathcal{E}_{\text{RPA}}^{\text{mass}} &= \frac{N_f^2 q^4 m}{48\pi^3} f(y) \quad \text{where } y = \left(\frac{16}{N_f}\right)^2 \frac{|B|}{q^3} \quad (\text{model C}) \\ f(y) &= y \int_0^\infty dk \frac{k^3(\sqrt{y}k + 1)}{(\sqrt{y}k + 1)^2(k^2 + 2)^2 + (16k^4/\pi^2)} \\ &\sim \frac{1}{2(1 + 16/\pi^2)} y \left\{ -\ln y + 1.8 \right\} \quad \text{for } y < 0.1 \quad . \end{aligned} \quad (16)$$

Notice that  $\Delta\mathcal{E}_{\text{RPA}}^{\text{mass}} > 0$ .

Secondly the lowest Landau level has an energy  $m_a$  so that a contribution from occupied levels is

$$\Delta\mathcal{E}_{\text{occupied}}^{\text{mass}} = \sum_{\text{light}} \frac{\nu_a m_a}{2\pi l_a^2} = \frac{1}{2\pi} \sum_{\text{light}} \nu_a m_a q_a |B| \quad . \quad (17)$$

It is positive.

There is another important contribution coming from a shift in fermion zero-point energies. Landau levels are given by  $\epsilon_{an} = \sqrt{m_a^2 + (2n/l_a^2)}$ . Recalling that there is asymmetry in the lowest level, i.e. the  $n = 0$  level exists for either a particle or an anti-particle,

one finds that fermion zero-point energies in the presence of a uniform magnetic field are

$$\Delta\mathcal{E}_{\text{f.z.e.}} = \sum_{\text{light}} \frac{1}{2\pi l_a^2} \left\{ -\frac{1}{2}m_a - \sum_{n=1}^{\infty} \sqrt{m_a^2 + \frac{2n}{l_a^2}} \right\} . \quad (18)$$

To evaluate this sum we utilize the standard zeta-function technique, taking

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^s} = \sum_n \frac{1}{\Gamma(s)} \int_0^{\infty} dt t^{s-1} e^{-\lambda_n t} . \quad (19)$$

Here we have  $\lambda_n \equiv m^2 + (2n/l^2)$  and need to calculate  $\zeta(-\frac{1}{2})$ . The substitution  $u \equiv 2nt/l^2$  immediately leads to

$$\begin{aligned} \zeta(s) &= \left(\frac{l^2}{2}\right)^s \frac{1}{\Gamma(s)} \sum_{n=1}^{\infty} \frac{1}{n^s} \int_0^{\infty} du u^{s-1} \exp\left(-u - \frac{m^2 l^2}{2n} u\right) \\ &= \left(\frac{l^2}{2}\right)^s \zeta_R(s) - m^2 \left(\frac{l^2}{2}\right)^{s+1} \zeta_R(s+1) + \cdots . \end{aligned} \quad (20)$$

Here  $\zeta_R$  is the Riemann zeta function and the expansion is valid for  $m^2 l^2 \ll 1$ . In particular

$$\zeta(-\frac{1}{2}) = -\frac{1}{4\pi} \zeta_R(\frac{3}{2}) \left(\frac{l^2}{2}\right)^{-1/2} - m^2 \left(\frac{l^2}{2}\right)^{1/2} \zeta_R(\frac{1}{2}) + \cdots . \quad (21)$$

Note  $\zeta_R(\frac{3}{2}) = 2.61238$  and  $\zeta_R(\frac{1}{2}) = -1.46035 < 0$ .

Utilizing (21) in (18), one finds

$$\begin{aligned} \Delta\mathcal{E}_{\text{f.z.e.}} &= \sum_{\text{light}} \frac{1}{4\sqrt{2}\pi^2} \zeta_R(\frac{3}{2}) \frac{1}{l_a^3} - \sum_{\text{light}} \frac{m_a}{4\pi l_a^2} + \mathcal{O}(m^2) \\ &= \frac{1}{4\sqrt{2}\pi^2} \zeta_R(\frac{3}{2}) \sum_{\text{light}} q_a^{3/2} |B|^{3/2} - \frac{1}{4\pi} \sum_{\text{light}} m_a q_a |B| + \mathcal{O}(m^2) \\ &\quad \text{for } \frac{m^2}{q^4} \ll \frac{|B|}{q^3} \ll 1 \end{aligned} \quad (22)$$

Notice that the second term (linear in  $m_a$ ) is negative. This behavior has been noticed by Gusynin, Miransky, and Shovkovy in a different context.[5] (17) and (22) combine to give

$$\begin{aligned} &\Delta\mathcal{E}_{\text{occupied}}^{\text{mass}} + \Delta\mathcal{E}_{\text{f.z.e.}} \\ &= \frac{1}{4\sqrt{2}\pi^2} \zeta_R(\frac{3}{2}) \sum_{\text{light}} q_a^{3/2} |B|^{3/2} + \frac{1}{2\pi} \sum_{\text{light}} (\nu_a - \frac{1}{2}) m_a q_a |B| + \mathcal{O}(m^2) . \end{aligned} \quad (23)$$



We recognize that in model B the second term vanishes.

Combining the results of (15), (16), and (23), one finds, in model C,

$$\begin{aligned}
\frac{1}{q^6} \Delta \mathcal{E} &= \frac{1}{2} \frac{B^2}{q^6} - \frac{N_f}{4\pi^3} \tan^{-1} \frac{4}{\pi} \cdot \frac{|B|}{q^3} + \frac{N_f \zeta_R(\frac{3}{2})}{4\sqrt{2}\pi^2} \cdot \left(\frac{|B|}{q^3}\right)^{3/2} \\
&\quad + \frac{N_f^2 m}{48\pi^3 q^2} f\left(\frac{256|B|}{N_f^2 q^3}\right) + \mathcal{O}(m^2, B^2) \\
&\sim \frac{1}{2} \left(\frac{N_f}{16}\right)^2 y^2 - 0.117 \left(\frac{N_f}{16}\right)^3 y + 0.749 \left(\frac{N_f}{16}\right)^4 y^{3/2} \\
&\quad + 0.0328 \left(\frac{N_f}{16}\right)^2 \frac{m}{q^2} y (-\ln y + 1.8) + \mathcal{O}(m^2) \\
&\quad \text{for } \frac{m^2}{q^4} \ll y = \left(\frac{16}{N_f}\right)^2 \frac{|B|}{q^3} < 0.1
\end{aligned} \tag{24}$$

We have restored the Maxwell energy term  $\frac{1}{2}B^2$ . Other  $\mathcal{O}(B^2)$  corrections have much smaller coefficients.

We observe that finite masses of light fermions tend to restore the Lorentz symmetry.  $\Delta \mathcal{E}(y)$  develops a minimum at  $y_{\min} \neq 0$ . As  $m$  gets bigger,  $y_{\min}$  decreases and  $\Delta \mathcal{E}(y_{\min})$  increases. More specifically

	$m/q^2$	$y_{\min}$	$\Delta \mathcal{E}_{\min}/q^6$	$(ml)^2 _{\min}$	
$N_f = 16$	0.	$9.2 \times 10^{-3}$	$-3.7 \times 10^{-4}$		(25)
	0.02	$8.7 \times 10^{-3}$	$-3.3 \times 10^{-4}$	0.05	
$N_f = 4$	0.	$1.0 \times 10^{-1}$	$-7.1 \times 10^{-5}$		
	0.02	$7.2 \times 10^{-2}$	$-3.5 \times 10^{-5}$	0.09	

For larger values of  $m/q^2$  the expressions (21) and (24) are not accurate.

In this paper we have shown that heavy fermions can induce spontaneous magnetization through the induced Chern-Simons coefficient at low energies. Further we showed that nonvanishing masses of light fermions tend to restore the Lorentz symmetry by giving a positive slope in the  $|B|$  dependence near the minimum of  $\Delta \mathcal{E}$ .

Our consideration here was limited to the dependence of the energy density on a dynamically generated magnetic field. In view of the recent result by Gusynin et al. [5] it

is of great interest to find the effective action as a function of both dynamically generated magnetic field  $B$  and chiral condensate  $\langle \bar{\psi}_a \psi_a \rangle$ . It may be that both  $B$  and  $\langle \bar{\psi}_a \psi_a \rangle$  are generated dynamically in a cooperative manner. We shall come back to this point in the near future.

*Note added:* Recently Kanemura and Matsushita have examined the behaviour of the model (1) at finite temperature in the massless fermion limit.[14] They have found that the coefficient of the linear term  $\propto |B|$  in the energy density (15) or (24) remains negative even at finite temperature.

### Acknowledgements

This work was supported in part by the U.S. Department of Energy under contract no. DE-AC02-83ER-40105. D.W. would like to thank C.L. Erickson and L.D. Reed for many stimulating and useful discussions.

## References

- [1] Y. Hosotani, Phys. Lett. 319B (1993) 332.
- [2] Y. Hosotani, Phys. Rev. D51 (1995) 2022.
- [3] P. Cea, Phys. Rev. D32 (1985) 2785.
- [4] Y. Hosotani, UMN-TH-1304/94 ([hep-th/9407188](#)); UMN-TH-1308/94 ([hep-th/9408148](#)).
- [5] V.P. Gusynin, V.A. Miransky, and I.A. Shovkovy, Phys. Rev. Lett. 73 (1994) 3499; NSF-ITP-94-74 ([hep-th/9407168](#)); Kiev-ITP-95-02E ([hep-ph/9501304](#)).
- [6] R.R. Parwani, IP/BBSR/95-12 ([hep-th/9504020](#)).
- [7] D. Cangemi, E. D'Hoker and G. Dunne, Phys. Rev. D51 (1995) 2513.
- [8] J.D. Bjorken, Ann. Phys. A30 (1963) 174.
- [9] Y. Nambu, Supplement of Prog. Theoret. Phys. Extra Number (1968) 190.
- [10] A. Kovner and B. Rosenstein, Int. J. Mod. Phys. A30 (1992) 7419.
- [11] A.N. Redlich, Phys. Rev. Lett. 52 (1984) 18; Phys. Rev. D29 (1984) 2366.

- [12] K. Ishikawa, Phys. Rev. Lett. 53 (1984) 1615; Phys. Rev. D31 (1985) 1432.
- [13] V.Y. Zeitlin, Yad. Fiz. 49 (1989) 742 [Sov. J. Nucl. Phys. 49 (1989) 461].
- [14] S. Kanemura and T. Matsushita, OU-HET 212 ([hep-th/9505146](#)) .